


# Diaquabis(2-oxo-2*H*-chromene-3-carboxylato- $\kappa^2 O^2, O^3$ )manganese(II)

Yue Cui, Qian Gao, Huan-Huan Wang, Lin Wang and Ya-Bo Xie

*Acta Cryst.* (2011). **E67**, m727

This open-access article is distributed under the terms of the Creative Commons Attribution Licence <http://creativecommons.org/licenses/by/2.0/uk/legalcode>, which permits unrestricted use, distribution, and reproduction in any medium, provided the original authors and source are cited.



*Acta Crystallographica Section E: Structure Reports Online* is the IUCr's highly popular open-access structural journal. It provides a simple and easily accessible publication mechanism for the growing number of inorganic, metal-organic and organic crystal structure determinations. The electronic submission, validation, refereeing and publication facilities of the journal ensure very rapid and high-quality publication, whilst key indicators and validation reports provide measures of structural reliability. The journal publishes over 4000 structures per year. The average publication time is less than one month.

Crystallography Journals **Online** is available from [journals.iucr.org](http://journals.iucr.org)

# Diaquabis(2-oxo-2H-chromene-3-carboxylato- $\kappa^2 O^2, O^3$ )manganese(II)

Yue Cui, Qian Gao, Huan-Huan Wang, Lin Wang and Ya-Bo Xie\*

College of Environmental and Energy Engineering, Beijing University of Technology, Beijing 100124, People's Republic of China  
Correspondence e-mail: xieyabo@bjut.edu.cn


Received 28 April 2011; accepted 3 May 2011

Key indicators: single-crystal X-ray study;  $T = 293$  K; mean  $\sigma(C-C) = 0.002$  Å;  $R$  factor = 0.027;  $wR$  factor = 0.070; data-to-parameter ratio = 13.8.

In the title compound,  $[Mn(C_{10}H_5O_4)_2(H_2O)_2]$ , the  $Mn^{II}$  atom lies on a crystallographic inversion center and is six-coordinated by two O atoms from water molecules in the axial positions and four O atoms from two deprotonated coumarin-3-carboxylic acid ligands in the equatorial plane. The overall coordination geometry is slightly distorted octahedral. The Mn—O bond distances vary between 2.0931 (12) and 2.2315 (13) Å. O—H...O hydrogen bonds between the H atoms of coordinated water molecules and the O atoms of the carboxylate groups link the complex molecules into two-dimensional layers parallel to the  $ab$  plane.

## Related literature

For background to topological networks, see: Hu *et al.* (2010). For applications of manganese(II) complexes, see: Hazra *et al.* (2011), Kuschel *et al.* (2010); Yang *et al.* (2010). For related structures, see: Gao *et al.* (2010).



## Experimental

### Crystal data

$[Mn(C_{10}H_5O_4)_2(H_2O)_2]$   $\alpha = 93.28$  (3)°  
 $M_r = 469.25$   $\beta = 90.67$  (3)°  
 Triclinic,  $P\bar{1}$   $\gamma = 113.47$  (3)°  
 $a = 6.7036$  (13) Å  $V = 446.33$  (15) Å<sup>3</sup>  
 $b = 6.9797$  (14) Å  $Z = 1$   
 $c = 10.424$  (2) Å Mo  $K\alpha$  radiation

$\mu = 0.80$  mm<sup>-1</sup>  
 $T = 293$  K

0.20 × 0.20 × 0.15 mm

### Data collection

Bruker APEXII CCD diffractometer  
 Absorption correction: multi-scan (SADABS; Bruker, 2008)  
 $T_{min} = 0.852$ ,  $T_{max} = 0.887$

2811 measured reflections  
 2021 independent reflections  
 1905 reflections with  $I > 2\sigma(I)$   
 $R_{int} = 0.009$

### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.027$   
 $wR(F^2) = 0.070$   
 $S = 1.07$   
 2021 reflections  
 146 parameters

H atoms treated by a mixture of independent and constrained refinement  
 $\Delta\rho_{max} = 0.27$  e Å<sup>-3</sup>  
 $\Delta\rho_{min} = -0.25$  e Å<sup>-3</sup>

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O1W—H1WA...O4 <sup>i</sup>	0.82	1.89	2.7113 (17)	178
O1W—H1WB...O4 <sup>ii</sup>	0.82 (3)	1.95 (3)	2.755 (2)	167 (2)

Symmetry codes: (i)  $-x+1, -y+1, -z+1$ ; (ii)  $x-1, y-1, z$ .

Data collection: APEX2 (Bruker, 2008); cell refinement: SAINT (Bruker, 2008); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: SHELXTL (Sheldrick, 2008); software used to prepare material for publication: SHELXL97.

This work was supported by the National Natural Science Foundation of China (No.21075114), the Science and Technology Development Project of the Beijing Education Committee and the Special Environmental Protection Fund for Public Welfare Project (201009015), the Funding Project for Academic Human Resources Development in Institutions of Higher Learning under the jurisdiction of the Beijing Municipality (PHR 201107104) and the Ninth Technology Fund for Postgraduates of Beijing University of Technology (ykj-2011-5406).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: SJ5131).

## References

- Bruker (2008). APEX2, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
- Gao, Q., Jiang, F. L., Wu, M. Y., Huang, Y. G., Wei, W. & Hong, M. C. (2010). *Cryst. Growth Des.*, **10**, 184–190.
- Hazra, A. J., Ali, Z. M., Shah, F. A., Zardari, L. A. & Khuhawar, M. Y. (2011). *Chem. Commun.* **47**, 538–540.
- Hu, J. S., Shang, Y. J., Yao, X. Q., Qin, L., Li, Y. Z., Guo, Z. J., Zheng, H. G. & Xue, Z. L. (2010). *Cryst. Growth Des.* **10**, 4135–4142.
- Kuschel, A., Luka, M., Wessig, M., Drescher, M., Fonin, M., Kiliani, G. & Polarz, S. (2010). *Adv. Funct. Mater.* **20**, 1133–1143.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Yang, Q., Zhao, J. P., Hu, B. W., Zhang, X. F. & Bu, X. H. (2010). *Inorg. Chem.* **49**, 3746–3751.

## **supplementary materials**

*Acta Cryst.* (2011). E67, m727 [ doi:10.1107/S1600536811016667 ]

## Diaquabis(2-oxo-2H-chromene-3-carboxylato- $\kappa^2 O^2, O^3$ )manganese(II)

Y. Cui, Q. Gao, H.-H. Wang, L. Wang and Y.-B. Xie

### Comment

In the past decades, numerous papers dealing with manganese(II) complexes have been published due to their fascinating structural diversity (Hu *et al.*, 2010) and potential applications in the areas of catalysis (Kuschel *et al.*, 2010), gas adsorption (Hazra *et al.*, 2011) and magnetism (Yang *et al.*, 2010). Herein, we report the synthesis and crystal structure of a new mononuclear manganese complex coordinated by coumarin-3-carboxylic acid.

In the title compound,  $[\text{Mn}(\text{C}_{10}\text{H}_5\text{O}_4)_2(\text{H}_2\text{O})_2]$ , each manganese(II) atom lies on a crystallographic inversion center and is six-coordinated by two O atoms from water molecules in the axial positions and four O atoms from two deprotonated coumarin-3-carboxylic acid ligands in the equatorial plane to exhibits a slightly distorted octahedral geometry. Angles around the Mn(II) atom vary between  $82.88(5)^\circ$  and  $97.12(5)^\circ$ . The Mn—O bond distances between the Mn(II) atom and the O atoms vary between 2.0931 (12) and 2.2315 (13) Å, all of which are comparable to those reported for other manganese-oxygen donor complexes (*e.g.* Gao *et al.*, 2010). The (C1/C2/C3/C4/C5/C6) and (C4/C3/C7/C8/C9/O1) rings are almost coplanar with a dihedral angle of  $1.847(6)^\circ$  between them. The dihedral angle between the C10/C8/C9/O2 and O2/Mn/O3 planes is  $25.803(7)^\circ$ . O—H $\cdots$ O hydrogen bonds between the hydrogen atoms of the coordinated water molecules and the O atoms of the carboxyl groups [O(1 W)—H(1 W A) $\cdots$ O(4), 2.711 (2) Å, O(1 W)—H(1WB) $\cdots$ O(4), 2.755 (2) Å] join the complexes into two-dimensional layers parallel the *ab* plane (Table 1, Fig. 2).

### Experimental

The title complex was synthesized by carefully layering a solution of  $\text{MnSO}_4 \cdot \text{H}_2\text{O}$  (16.9 mg, 0.1 mmol) in ethanol (10 ml) on top of a solution of coumarin-3-carboxylic acid (19.0 mg, 0.1 mmol) and LiOH (8.4 mg, 0.2 mmol) in  $\text{H}_2\text{O}$  (10 ml) in a test-tube. After about one month at room temperature, yellow block-shaped single crystals suitable for X-ray investigation appeared at the boundary between the ethanol solution and the water layer with a yield of 25% (12.1 mg). FT—IR (KBr,  $\text{cm}^{-1}$ ): 788, 1028, 1183, 1285, 1388, 1457, 1585, 1662, 3214.

### Refinement

Carbon H atoms were placed geometrically (C—H = 0.93 Å) and treated as riding with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ . Water H atoms were located in a difference Fourier map. One was treated as riding in the subsequent refinement atoms, with O—H = 0.85 Å, the coordinates of the other were refined freely. For both  $i > U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{O})$ .

## Figures




Fig. 1. The molecular structure of the title compound with displacement ellipsoids drawn at the 50% probability level for non-hydrogen atoms, hydrogen atoms are shown as small circles of arbitrary radius. [Symmetry code: A = -x, -y + 1, -z + 1].




Fig. 2. Partial packing view of title compound, showing the formation of network built from hydrogen bonds.

## Diaquabis(2-oxo-2H-chromene-3-carboxylato- $\kappa^2 O^2, O^3$ )manganese(II)

### Crystal data

[Mn(C<sub>10</sub>H<sub>5</sub>O<sub>4</sub>)<sub>2</sub>(H<sub>2</sub>O)<sub>2</sub>]

$M_r = 469.25$

Triclinic,  $P\bar{1}$

Hall symbol: -P 1

$a = 6.7036$  (13) Å

$b = 6.9797$  (14) Å

$c = 10.424$  (2) Å

$\alpha = 93.28$  (3)°

$\beta = 90.67$  (3)°

$\gamma = 113.47$  (3)°

$V = 446.33$  (15) Å<sup>3</sup>

$Z = 1$

$F(000) = 239$

$D_x = 1.746$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 1947 reflections

$\theta = 2.0$ – $28.3^\circ$

$\mu = 0.80$  mm<sup>-1</sup>

$T = 293$  K

Block, yellow

$0.20 \times 0.20 \times 0.15$  mm

### Data collection

Bruker APEXII CCD  
diffractometer

Radiation source: fine-focus sealed tube  
graphite

$\varphi$  and  $\omega$  scans

Absorption correction: multi-scan  
(*SADABS*; Bruker, 2008)

$T_{\min} = 0.852$ ,  $T_{\max} = 0.887$

2811 measured reflections

2021 independent reflections

1905 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.009$

$\theta_{\max} = 28.3^\circ$ ,  $\theta_{\min} = 2.0^\circ$

$h = -7 \rightarrow 8$

$k = -8 \rightarrow 8$

$l = -13 \rightarrow 13$

### Refinement

Refinement on  $F^2$

Primary atom site location: structure-invariant direct  
methods

Least-squares matrix: full

$$R[F^2 > 2\sigma(F^2)] = 0.027$$

$$wR(F^2) = 0.070$$

$$S = 1.07$$

2021 reflections

146 parameters

0 restraints

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of independent and constrained refinement

$$w = 1/[\sigma^2(F_o^2) + (0.030P)^2 + 0.2201P]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} = 0.002$$

$$\Delta\rho_{\max} = 0.27 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.25 \text{ e } \text{\AA}^{-3}$$

### Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Mn1	0.0000	0.5000	0.5000	0.02451 (10)
O1	0.20420 (17)	0.69142 (18)	0.12393 (10)	0.0279 (2)
O1W	−0.00060 (19)	0.21428 (19)	0.40159 (11)	0.0308 (2)
H1WA	0.0974	0.1872	0.4325	0.046*
O2	0.06195 (17)	0.64929 (19)	0.31264 (11)	0.0311 (3)
O3	0.34009 (16)	0.63381 (18)	0.51888 (10)	0.0290 (2)
O4	0.67461 (16)	0.86812 (18)	0.49213 (11)	0.0307 (3)
C1	0.7033 (3)	0.7877 (3)	−0.12676 (17)	0.0377 (4)
H1A	0.8140	0.8088	−0.1842	0.045*
C2	0.7458 (3)	0.7881 (3)	0.00281 (16)	0.0327 (3)
H2A	0.8845	0.8080	0.0326	0.039*
C3	0.5793 (2)	0.7581 (2)	0.09021 (14)	0.0255 (3)
C4	0.3735 (2)	0.7261 (2)	0.04195 (14)	0.0254 (3)
C5	0.3284 (3)	0.7240 (3)	−0.08835 (15)	0.0332 (4)
H5A	0.1893	0.7017	−0.1186	0.040*
C6	0.4953 (3)	0.7559 (3)	−0.17236 (16)	0.0377 (4)
H6A	0.4685	0.7562	−0.2601	0.045*
C7	0.6108 (2)	0.7651 (2)	0.22644 (14)	0.0253 (3)
H7A	0.7483	0.7901	0.2607	0.030*
C8	0.4463 (2)	0.7363 (2)	0.30689 (13)	0.0218 (3)
C9	0.2290 (2)	0.6896 (2)	0.25373 (14)	0.0230 (3)
C10	0.4883 (2)	0.7473 (2)	0.45064 (14)	0.0223 (3)

# supplementary materials

H1WB                      −0.110 (4)                      0.112 (4)                      0.418 (2)                      0.057 (7)\*

## Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Mn1	0.01647 (15)	0.02948 (18)	0.02242 (17)	0.00348 (12)	0.00291 (11)	0.00327 (12)
O1	0.0219 (5)	0.0382 (6)	0.0210 (5)	0.0088 (4)	−0.0001 (4)	0.0052 (4)
O1W	0.0233 (5)	0.0326 (6)	0.0332 (6)	0.0077 (5)	0.0017 (4)	0.0022 (5)
O2	0.0191 (5)	0.0457 (7)	0.0274 (6)	0.0107 (5)	0.0038 (4)	0.0102 (5)
O3	0.0188 (5)	0.0397 (6)	0.0220 (5)	0.0042 (4)	0.0017 (4)	0.0066 (4)
O4	0.0185 (5)	0.0351 (6)	0.0297 (6)	0.0014 (4)	−0.0035 (4)	0.0037 (5)
C1	0.0473 (10)	0.0351 (9)	0.0282 (8)	0.0133 (7)	0.0157 (7)	0.0038 (7)
C2	0.0310 (8)	0.0342 (8)	0.0307 (8)	0.0104 (6)	0.0093 (6)	0.0030 (6)
C3	0.0263 (7)	0.0253 (7)	0.0223 (7)	0.0073 (6)	0.0042 (5)	0.0030 (5)
C4	0.0271 (7)	0.0236 (7)	0.0223 (7)	0.0066 (6)	0.0033 (5)	0.0030 (5)
C5	0.0389 (9)	0.0324 (8)	0.0242 (8)	0.0101 (7)	−0.0031 (6)	0.0016 (6)
C6	0.0558 (11)	0.0332 (8)	0.0199 (7)	0.0133 (8)	0.0046 (7)	0.0020 (6)
C7	0.0202 (6)	0.0277 (7)	0.0257 (7)	0.0072 (5)	0.0007 (5)	0.0020 (6)
C8	0.0186 (6)	0.0237 (7)	0.0210 (7)	0.0060 (5)	0.0001 (5)	0.0022 (5)
C9	0.0217 (7)	0.0241 (7)	0.0213 (7)	0.0066 (5)	0.0009 (5)	0.0049 (5)
C10	0.0174 (6)	0.0258 (7)	0.0224 (7)	0.0074 (5)	0.0000 (5)	0.0016 (5)

## Geometric parameters ( $\text{\AA}$ , $^\circ$ )

Mn1—O3	2.0931 (12)	C1—C6	1.395 (3)
Mn1—O3 <sup>i</sup>	2.0931 (12)	C1—H1A	0.9300
Mn1—O1W	2.1884 (13)	C2—C3	1.407 (2)
Mn1—O1W <sup>i</sup>	2.1884 (13)	C2—H2A	0.9300
Mn1—O2 <sup>i</sup>	2.2315 (13)	C3—C4	1.389 (2)
Mn1—O2	2.2315 (13)	C3—C7	1.429 (2)
O1—C9	1.3626 (17)	C4—C5	1.386 (2)
O1—C4	1.3802 (18)	C5—C6	1.384 (3)
O1W—H1WA	0.8200	C5—H5A	0.9300
O1W—H1WB	0.82 (3)	C6—H6A	0.9300
O2—C9	1.2225 (18)	C7—C8	1.351 (2)
O3—C10	1.2549 (18)	C7—H7A	0.9300
O4—C10	1.2509 (17)	C8—C9	1.4555 (19)
C1—C2	1.377 (2)	C8—C10	1.5133 (19)
O3—Mn1—O3 <sup>i</sup>	180.0	C3—C2—H2A	120.0
O3—Mn1—O1W	91.96 (5)	C4—C3—C2	118.33 (14)
O3 <sup>i</sup> —Mn1—O1W	88.04 (5)	C4—C3—C7	117.86 (14)
O3—Mn1—O1W <sup>i</sup>	88.04 (5)	C2—C3—C7	123.78 (14)
O3 <sup>i</sup> —Mn1—O1W <sup>i</sup>	91.96 (5)	O1—C4—C5	117.46 (14)
O1W—Mn1—O1W <sup>i</sup>	180.00 (6)	O1—C4—C3	120.28 (13)
O3—Mn1—O2 <sup>i</sup>	97.12 (5)	C5—C4—C3	122.25 (15)
O3 <sup>i</sup> —Mn1—O2 <sup>i</sup>	82.88 (5)	C4—C5—C6	118.39 (16)

O1W—Mn1—O2 <sup>i</sup>	91.16 (5)	C4—C5—H5A	120.8
O1W <sup>i</sup> —Mn1—O2 <sup>i</sup>	88.84 (5)	C6—C5—H5A	120.8
O3—Mn1—O2	82.88 (5)	C5—C6—C1	120.65 (16)
O3 <sup>i</sup> —Mn1—O2	97.12 (5)	C5—C6—H6A	119.7
O1W—Mn1—O2	88.84 (5)	C1—C6—H6A	119.7
O1W <sup>i</sup> —Mn1—O2	91.16 (5)	C8—C7—C3	121.79 (14)
O2 <sup>i</sup> —Mn1—O2	180.0	C8—C7—H7A	119.1
C9—O1—C4	122.73 (12)	C3—C7—H7A	119.1
Mn1—O1W—H1WA	109.5	C7—C8—C9	119.37 (13)
Mn1—O1W—H1WB	110.8 (17)	C7—C8—C10	119.84 (13)
H1WA—O1W—H1WB	102.0	C9—C8—C10	120.77 (12)
C9—O2—Mn1	124.45 (10)	O2—C9—O1	114.86 (13)
C10—O3—Mn1	133.91 (10)	O2—C9—C8	127.35 (14)
C2—C1—C6	120.41 (16)	O1—C9—C8	117.78 (13)
C2—C1—H1A	119.8	O4—C10—O3	124.70 (14)
C6—C1—H1A	119.8	O4—C10—C8	116.15 (13)
C1—C2—C3	119.96 (16)	O3—C10—C8	119.13 (12)
C1—C2—H2A	120.0		
O3—Mn1—O2—C9	25.72 (12)	C2—C1—C6—C5	−0.1 (3)
O3 <sup>i</sup> —Mn1—O2—C9	−154.28 (12)	C4—C3—C7—C8	1.7 (2)
O1W—Mn1—O2—C9	−66.40 (13)	C2—C3—C7—C8	179.71 (15)
O1W <sup>i</sup> —Mn1—O2—C9	113.60 (13)	C3—C7—C8—C9	2.2 (2)
O1W—Mn1—O3—C10	90.62 (15)	C3—C7—C8—C10	−179.37 (13)
O1W <sup>i</sup> —Mn1—O3—C10	−89.38 (15)	Mn1—O2—C9—O1	152.34 (10)
O2 <sup>i</sup> —Mn1—O3—C10	−177.97 (14)	Mn1—O2—C9—C8	−28.2 (2)
O2—Mn1—O3—C10	2.03 (14)	C4—O1—C9—O2	−177.49 (13)
C6—C1—C2—C3	−0.6 (3)	C4—O1—C9—C8	3.0 (2)
C1—C2—C3—C4	0.8 (2)	C7—C8—C9—O2	176.00 (15)
C1—C2—C3—C7	−177.20 (15)	C10—C8—C9—O2	−2.4 (2)
C9—O1—C4—C5	179.93 (14)	C7—C8—C9—O1	−4.5 (2)
C9—O1—C4—C3	1.0 (2)	C10—C8—C9—O1	177.09 (12)
C2—C3—C4—O1	178.54 (13)	Mn1—O3—C10—O4	155.33 (12)
C7—C3—C4—O1	−3.3 (2)	Mn1—O3—C10—C8	−26.6 (2)
C2—C3—C4—C5	−0.4 (2)	C7—C8—C10—O4	31.1 (2)
C7—C3—C4—C5	177.76 (14)	C9—C8—C10—O4	−150.50 (14)
O1—C4—C5—C6	−179.23 (14)	C7—C8—C10—O3	−147.10 (15)
C3—C4—C5—C6	−0.3 (2)	C9—C8—C10—O3	31.3 (2)
C4—C5—C6—C1	0.5 (3)		

Symmetry codes: (i)  $-x, -y+1, -z+1$ .

### Hydrogen-bond geometry ( $\text{\AA}$ , $^\circ$ )

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O1W—H1WA $\cdots$ O4 <sup>ii</sup>	0.82	1.89	2.7113 (17)	178
O1W—H1WB $\cdots$ O4 <sup>iii</sup>	0.82 (3)	1.95 (3)	2.755 (2)	167 (2)

Symmetry codes: (ii)  $-x+1, -y+1, -z+1$ ; (iii)  $x-1, y-1, z$ .



Fig. 1




Fig. 2

