


# Diaquabis(2-oxo-2*H*-chromene-3-carboxylato- $\kappa^2 O^2, O^3$ )cadmium

Yue Cui, Qian Gao, Huan-Huan Wang, Lin Wang and Ya-Bo Xie

*Acta Cryst.* (2011). **E67**, m126

This open-access article is distributed under the terms of the Creative Commons Attribution Licence <http://creativecommons.org/licenses/by/2.0/uk/legalcode>, which permits unrestricted use, distribution, and reproduction in any medium, provided the original authors and source are cited.



*Acta Crystallographica Section E: Structure Reports Online* is the IUCr's highly popular open-access structural journal. It provides a simple and easily accessible publication mechanism for the growing number of inorganic, metal-organic and organic crystal structure determinations. The electronic submission, validation, refereeing and publication facilities of the journal ensure very rapid and high-quality publication, whilst key indicators and validation reports provide measures of structural reliability. In 2007, the journal published over 5000 structures. The average publication time is less than one month.

Crystallography Journals **Online** is available from [journals.iucr.org](http://journals.iucr.org)

## Diaquabis(2-oxo-2H-chromene-3-carboxylato- $\kappa^2 O^2, O^3$ )cadmium

Yue Cui, Qian Gao, Huan-Huan Wang, Lin Wang and Ya-Bo Xie\*

College of Environmental and Energy Engineering, Beijing University of Technology, Beijing 100124, People's Republic of China  
Correspondence e-mail: xieyabo@bjut.edu.cn


Received 1 December 2010; accepted 20 December 2010

Key indicators: single-crystal X-ray study;  $T = 110$  K; mean  $\sigma(C-C) = 0.003$  Å;  $R$  factor = 0.017;  $wR$  factor = 0.045; data-to-parameter ratio = 14.4.

In the title mononuclear cadmium complex,  $[Cd(C_{10}H_5O_4)_2 \cdot (H_2O)_2]$ , the  $Cd^{II}$  atom, located on a crystallographic inversion center, exhibits a slightly distorted octahedral geometry and is six-coordinated by two O atoms from water molecules in the axial positions and four O atoms from two deprotonated coumarin-3-carboxylic acid ligands in the equatorial plane. Angles around the  $Cd^{II}$  atom vary between  $81.00$  (5) and  $99.00$  (0)°. The  $Cd-O$  bond lengths vary between  $2.1961$  (13) and  $2.3360$  (13) Å.  $O-H \cdots O$  hydrogen bonds between the H atoms of coordinated water molecules and the O atoms of carboxylate groups link the complex molecules into layers parallel to the  $ab$  plane.

### Related literature

For background to topological networks, see: Lin *et al.* (2010). For applications of self-assembling systems with organic ligands containing O donors, see: Bischof *et al.* (2010); Chen *et al.* (2008); Ghoshal *et al.* (2007); Li & Zhou (2009). For related structures, see: Georgieva *et al.* (2007); Li *et al.* (2009).



### Experimental

#### Crystal data

$[Cd(C_{10}H_5O_4)_2 \cdot (H_2O)_2]$   
 $M_r = 526.72$   
Triclinic,  $P\bar{1}$   
 $a = 6.6736$  (13) Å  
 $b = 6.8838$  (14) Å  
 $c = 10.477$  (2) Å  
 $\alpha = 93.37$  (3)°  
 $\beta = 91.46$  (3)°

$\gamma = 112.07$  (3)°  
 $V = 444.68$  (15) Å<sup>3</sup>  
 $Z = 1$   
Mo  $K\alpha$  radiation

$\mu = 1.29$  mm<sup>-1</sup>  
 $T = 110$  K  
 $0.20 \times 0.15 \times 0.15$  mm

#### Data collection

Bruker APEXII CCD diffractometer  
Absorption correction: multi-scan (SADABS; Sheldrick, 2008a)  
 $T_{min} = 0.793$ ,  $T_{max} = 0.824$

2812 measured reflections  
2040 independent reflections  
2033 reflections with  $I > 2\sigma(I)$   
 $R_{int} = 0.009$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.017$   
 $wR(F^2) = 0.045$   
 $S = 1.12$   
2040 reflections

142 parameters  
H-atom parameters constrained  
 $\Delta\rho_{max} = 0.42$  e Å<sup>-3</sup>  
 $\Delta\rho_{min} = -0.44$  e Å<sup>-3</sup>

**Table 1**

Hydrogen-bond geometry (Å, °).

| $D-H \cdots A$            | $D-H$ | $H \cdots A$ | $D \cdots A$ | $D-H \cdots A$ |
|---------------------------|-------|--------------|--------------|----------------|
| $O1W-H1WA \cdots O4^i$    | 0.85  | 1.90         | 2.6877 (18)  | 153            |
| $O1W-H1WB \cdots O4^{ii}$ | 0.85  | 1.94         | 2.721 (2)    | 153            |

Symmetry codes: (i)  $-x+1, -y+1, -z$ ; (ii)  $x+1, y+1, z$ .

Data collection: APEX2 (Bruker, 2008); cell refinement: SAINT (Bruker, 2008); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008b); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008b); molecular graphics: SHELXTL (Sheldrick, 2008b); software used to prepare material for publication: SHELXL97.

This work was supported by the National Natural Science Foundation of China (No. 21075114), the Science and Technology Development Project of Beijing Education Committee and the Special Environmental Protection Fund for Public Welfare project (201009015).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: ZL2335).

### References

- Bischof, S. M., Ess, D. H., Meier, S. K., Oxgaard, J., Nielsen, R. J., Bhalla, G., Goddard, W. A. & Periana, R. A. (2010). *Organometallics*, **29**, 742–756.
- Bruker (2008). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
- Chen, L. F., Li, Z. J., Qin, Y. Y., Cheng, J. K. & Yao, Y. G. (2008). *J. Mol. Struct.* **892**, 278–282.
- Georgieva, I., Trendafilova, N., Aquino, A. J. A. & Lischka, H. (2007). *Inorg. Chem.* **46**, 10926–10936.
- Ghoshal, D., Ghosh, A. K., Mostafa, G., Ribas, J. & Chaudhuri, N. R. (2007). *Inorg. Chim. Acta*, **360**, 1771–1775.
- Li, N., Gou, L., Hu, H. M., Chen, S. H., Chen, X. L., Wang, B. W., Wu, Q. R., Yang, M. L. & Xue, G. L. (2009). *Inorg. Chim. Acta*, **362**, 3475–3483.
- Li, J. R. & Zhou, H. C. (2009). *Angew. Chem. Int. Ed.* **48**, 1–5.
- Lin, J. D., Long, X. F., Lin, P. & Du, S. W. (2010). *Cryst. Growth Des.* **10**, 146–157.
- Sheldrick, G. M. (2008a). SADABS. University of Göttingen, Germany.
- Sheldrick, G. M. (2008b). *Acta Cryst.* **A64**, 112–122.

## **supplementary materials**

*Acta Cryst.* (2011). E67, m126 [ doi:10.1107/S1600536810053523 ]

## Diaquabis(2-oxo-2H-chromene-3-carboxylato- $\kappa^2 O^2, O^3$ )cadmium

Y. Cui, Q. Gao, H.-H. Wang, L. Wang and Y.-B. Xie

### Comment

In the past decades, much attention has been paid to the design and synthesis of self-assembling metal complex systems with organic ligands containing O donors due to their fascinating structural diversity (Lin *et al.*, 2010) and potential applications in the areas of catalysis (Bischof *et al.*, 2010), magnetism (Ghoshal *et al.*, 2007), gas adsorption (Li & Zhou, 2009), and luminescence (Chen *et al.*, 2008). Coumarin-3-carboxylic acid is such a ligand and complexes containing it have been reported (Georgieva *et al.*, 2007). Herein, we report the synthesis and crystal structure of a new mononuclear cadmium complex coordinated by coumarin-3-carboxylic acid.

The molecule of the title mononuclear cadmium(II) complex,  $[\text{Cd}(\text{C}_{10}\text{H}_5\text{O}_4)_2(\text{H}_2\text{O})_2]$ , occupies a special position with the metal center being located on a crystallographic inversion center. Each  $\text{Cd}^{\text{II}}$  atom exhibits a slightly distorted octahedral geometry and is six-coordinated by two O atoms from water molecules in the axial positions and four O atoms from two deprotonated coumarin-3-carboxylic acid ligands in the equatorial plane. Angles around the  $\text{Cd}^{\text{II}}$  atom vary between  $81.02(6)^\circ$  and  $98.98(8)^\circ$ . The Cd—O bond distances between the  $\text{Cd}^{\text{II}}$  atom and the O atoms vary between 2.196(2) and 2.336(2) Å, all of which are comparable to those reported for other cadmium-oxygen donor complexes (e.g., Li *et al.*, 2009). The (C1C2C3C4C5C6) ring and the (C6C5C7C8C9O1) ring are almost coplanar, and the dihedral angle is  $1.673(5)^\circ$ . The dihedral angle between the C8C9C10O2 plane and the O2O3Cd1 plane is  $28.541(7)^\circ$ . O—H...O hydrogen bonds between the hydrogen atoms of coordinated water molecules and the O atoms of carboxyl groups joins the complexes into two-dimensional layers parallel the *ab* plane (Table 1, Fig. 2).

### Experimental

The title complex was synthesized by carefully layering a solution of  $\text{Cd}(\text{NO}_3)_2 \cdot 4\text{H}_2\text{O}$  (30.8 mg, 0.1 mmol) in ethanol (10 ml) on top of a solution of coumarin-3-carboxylic acid (19.0 mg, 0.1 mmol) and LiOH (8.4 mg, 0.2 mmol) in  $\text{H}_2\text{O}$  (10 ml) in a test-tube. After about one month at room temperature, colorless block-shaped single crystals suitable for X-ray investigation appeared at the boundary between the ethanol solution and the water layer with a yield of 23% (12.1 mg). Decomposition temperature: near 573K. FT-IR (KBr,  $\text{cm}^{-1}$ ): 636, 748, 777, 1183, 1281, 1394, 1450, 1562, 1604, 1674.

### Refinement

Carbon H atoms were placed geometrically (C—H = 0.93 Å) and treated as riding with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ . Water H atoms were located in calculated positions and treated in the subsequent refinement as riding atoms, with O—H = 0.85 Å and  $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{O})$ .

## Figures




Fig. 1. The molecular structure of the title compound with displacement ellipsoids drawn at the 50% probability level for non-hydrogen atoms, hydrogen atoms are shown as small circles of arbitrary radius. [Symmetry code:  $i = -x + 2, -y + 1, -z$ ].




Fig. 2. Partial packing view of title compound, showing the formation of network built from hydrogen bonds.

## Diaquabis(2-oxo-2H-chromene-3-carboxylato- $\kappa^2 O^2, O^3$ )cadmium

### Crystal data

$[\text{Cd}(\text{C}_{10}\text{H}_5\text{O}_4)_2(\text{H}_2\text{O})_2]$

$M_r = 526.72$

Triclinic,  $P\bar{1}$

Hall symbol:  $-P\ 1$

$a = 6.6736\ (13)\ \text{\AA}$

$b = 6.8838\ (14)\ \text{\AA}$

$c = 10.477\ (2)\ \text{\AA}$

$\alpha = 93.37\ (3)^\circ$

$\beta = 91.46\ (3)^\circ$

$\gamma = 112.07\ (3)^\circ$

$V = 444.68\ (15)\ \text{\AA}^3$

$Z = 1$

$F(000) = 262$

$D_x = 1.967\ \text{Mg m}^{-3}$

Mo  $K\alpha$  radiation,  $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 2723 reflections

$\theta = 3.2\text{--}28.3^\circ$

$\mu = 1.29\ \text{mm}^{-1}$

$T = 110\ \text{K}$

Block, colorless

$0.20 \times 0.15 \times 0.15\ \text{mm}$

### Data collection

Bruker APEXII CCD  
diffractometer

Radiation source: fine-focus sealed tube  
graphite

$\varphi$  and  $\omega$  scans

Absorption correction: multi-scan  
(*SADABS*; Sheldrick, 2008a)

$T_{\min} = 0.793$ ,  $T_{\max} = 0.824$

2812 measured reflections

2040 independent reflections

2033 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.009$

$\theta_{\max} = 28.3^\circ$ ,  $\theta_{\min} = 3.2^\circ$

$h = -8 \rightarrow 8$

$k = -8 \rightarrow 8$

$l = -13 \rightarrow 6$

### Refinement

Refinement on  $F^2$

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.017$

$wR(F^2) = 0.045$

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

|                  |                                                        |
|------------------|--------------------------------------------------------|
| $S = 1.12$       | $w = 1/[\sigma^2(F_o^2) + (0.0217P)^2 + 0.3208P]$      |
| 2040 reflections | where $P = (F_o^2 + 2F_c^2)/3$                         |
| 142 parameters   | $(\Delta/\sigma)_{\max} < 0.001$                       |
| 0 restraints     | $\Delta\rho_{\max} = 0.42 \text{ e } \text{\AA}^{-3}$  |
|                  | $\Delta\rho_{\min} = -0.44 \text{ e } \text{\AA}^{-3}$ |

### Special details

**Geometry.** All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted R-factor wR and goodness of fit S are based on  $F^2$ , conventional R-factors R are based on F, with F set to zero for negative  $F^2$ . The threshold expression of  $F^2 > 2\text{sigma}(F^2)$  is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on  $F^2$  are statistically about twice as large as those based on F, and R- factors based on ALL data will be even larger.

### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

|      | x            | y            | z             | $U_{\text{iso}}^*/U_{\text{eq}}$ |
|------|--------------|--------------|---------------|----------------------------------|
| Cd1  | 1.0000       | 0.5000       | 0.0000        | 0.01268 (6)                      |
| O1   | 0.79598 (18) | 0.30467 (18) | 0.37932 (11)  | 0.0144 (2)                       |
| O1W  | 1.00099 (19) | 0.79679 (19) | 0.10974 (11)  | 0.0163 (2)                       |
| H1WA | 0.8758       | 0.7921       | 0.0875        | 0.024*                           |
| H1WB | 1.1256       | 0.8837       | 0.0939        | 0.024*                           |
| O2   | 0.93465 (18) | 0.33456 (19) | 0.19160 (11)  | 0.0165 (2)                       |
| O3   | 0.64486 (18) | 0.35849 (19) | -0.01845 (11) | 0.0161 (2)                       |
| O4   | 0.31702 (18) | 0.12595 (19) | 0.01275 (11)  | 0.0164 (2)                       |
| C1   | 0.6758 (3)   | 0.2758 (3)   | 0.59042 (16)  | 0.0170 (3)                       |
| H1A  | 0.8180       | 0.2975       | 0.6218        | 0.020*                           |
| C2   | 0.5088 (3)   | 0.2453 (3)   | 0.67327 (16)  | 0.0186 (3)                       |
| H2A  | 0.5371       | 0.2462       | 0.7627        | 0.022*                           |
| C3   | 0.2996 (3)   | 0.2132 (3)   | 0.62646 (16)  | 0.0192 (3)                       |
| H3A  | 0.1869       | 0.1917       | 0.6841        | 0.023*                           |
| C4   | 0.2562 (3)   | 0.2127 (3)   | 0.49686 (16)  | 0.0162 (3)                       |
| H4A  | 0.1144       | 0.1926       | 0.4657        | 0.019*                           |
| C5   | 0.4221 (3)   | 0.2421 (2)   | 0.41095 (15)  | 0.0137 (3)                       |
| C6   | 0.6282 (3)   | 0.2734 (2)   | 0.46024 (15)  | 0.0138 (3)                       |
| C7   | 0.3889 (3)   | 0.2351 (2)   | 0.27516 (15)  | 0.0131 (3)                       |
| H7A  | 0.2487       | 0.2123       | 0.2398        | 0.016*                           |
| C8   | 0.5526 (3)   | 0.2604 (2)   | 0.19545 (15)  | 0.0123 (3)                       |
| C9   | 0.7689 (3)   | 0.3026 (2)   | 0.24937 (15)  | 0.0127 (3)                       |
| C10  | 0.5040 (3)   | 0.2474 (2)   | 0.05228 (15)  | 0.0127 (3)                       |

## Atomic displacement parameters ( $\text{\AA}^2$ )

|     | $U^{11}$    | $U^{22}$    | $U^{33}$    | $U^{12}$    | $U^{13}$    | $U^{23}$    |
|-----|-------------|-------------|-------------|-------------|-------------|-------------|
| Cd1 | 0.00892 (8) | 0.01495 (9) | 0.01174 (9) | 0.00156 (6) | 0.00160 (5) | 0.00172 (5) |
| O1  | 0.0121 (5)  | 0.0186 (6)  | 0.0118 (5)  | 0.0048 (4)  | 0.0005 (4)  | 0.0029 (4)  |
| O1W | 0.0119 (5)  | 0.0180 (6)  | 0.0174 (6)  | 0.0038 (4)  | 0.0014 (4)  | 0.0014 (4)  |
| O2  | 0.0116 (5)  | 0.0228 (6)  | 0.0155 (6)  | 0.0062 (5)  | 0.0023 (4)  | 0.0057 (4)  |
| O3  | 0.0111 (5)  | 0.0209 (6)  | 0.0138 (5)  | 0.0027 (5)  | 0.0006 (4)  | 0.0037 (4)  |
| O4  | 0.0109 (5)  | 0.0179 (6)  | 0.0164 (5)  | 0.0010 (4)  | −0.0018 (4) | 0.0020 (4)  |
| C1  | 0.0190 (8)  | 0.0147 (7)  | 0.0158 (8)  | 0.0048 (6)  | −0.0011 (6) | 0.0018 (6)  |
| C2  | 0.0285 (9)  | 0.0148 (7)  | 0.0119 (7)  | 0.0072 (7)  | 0.0018 (6)  | 0.0015 (6)  |
| C3  | 0.0242 (9)  | 0.0162 (8)  | 0.0168 (8)  | 0.0068 (7)  | 0.0083 (6)  | 0.0019 (6)  |
| C4  | 0.0158 (7)  | 0.0147 (7)  | 0.0176 (8)  | 0.0048 (6)  | 0.0036 (6)  | 0.0021 (6)  |
| C5  | 0.0155 (7)  | 0.0107 (7)  | 0.0140 (7)  | 0.0037 (6)  | 0.0016 (6)  | 0.0012 (5)  |
| C6  | 0.0155 (7)  | 0.0108 (7)  | 0.0143 (7)  | 0.0037 (6)  | 0.0036 (6)  | 0.0018 (5)  |
| C7  | 0.0116 (7)  | 0.0126 (7)  | 0.0147 (7)  | 0.0041 (6)  | 0.0005 (6)  | 0.0020 (5)  |
| C8  | 0.0119 (7)  | 0.0117 (7)  | 0.0129 (7)  | 0.0037 (6)  | 0.0005 (5)  | 0.0018 (5)  |
| C9  | 0.0132 (7)  | 0.0118 (7)  | 0.0124 (7)  | 0.0037 (6)  | −0.0005 (5) | 0.0020 (5)  |
| C10 | 0.0114 (7)  | 0.0138 (7)  | 0.0133 (7)  | 0.0055 (6)  | −0.0002 (5) | 0.0006 (5)  |

## Geometric parameters ( $\text{\AA}$ , $^\circ$ )

|                                       |             |           |             |
|---------------------------------------|-------------|-----------|-------------|
| Cd1—O3 <sup>i</sup>                   | 2.1961 (13) | C1—C2     | 1.391 (2)   |
| Cd1—O3                                | 2.1961 (13) | C1—H1A    | 0.9500      |
| Cd1—O1W                               | 2.2824 (13) | C2—C3     | 1.400 (3)   |
| Cd1—O1W <sup>i</sup>                  | 2.2824 (13) | C2—H2A    | 0.9500      |
| Cd1—O2 <sup>i</sup>                   | 2.3360 (13) | C3—C4     | 1.381 (2)   |
| Cd1—O2                                | 2.3360 (13) | C3—H3A    | 0.9500      |
| O1—C9                                 | 1.3673 (19) | C4—C5     | 1.407 (2)   |
| O1—C6                                 | 1.3810 (19) | C4—H4A    | 0.9500      |
| O1W—H1WA                              | 0.8500      | C5—C6     | 1.390 (2)   |
| O1W—H1WB                              | 0.8500      | C5—C7     | 1.430 (2)   |
| O2—C9                                 | 1.227 (2)   | C7—C8     | 1.357 (2)   |
| O3—C10                                | 1.257 (2)   | C7—H7A    | 0.9500      |
| O4—C10                                | 1.256 (2)   | C8—C9     | 1.453 (2)   |
| C1—C6                                 | 1.390 (2)   | C8—C10    | 1.517 (2)   |
| O3 <sup>i</sup> —Cd1—O3               | 180.0       | C3—C2—H2A | 119.6       |
| O3 <sup>i</sup> —Cd1—O1W              | 87.17 (5)   | C4—C3—C2  | 120.30 (16) |
| O3—Cd1—O1W                            | 92.83 (5)   | C4—C3—H3A | 119.8       |
| O3 <sup>i</sup> —Cd1—O1W <sup>i</sup> | 92.83 (5)   | C2—C3—H3A | 119.8       |
| O3—Cd1—O1W <sup>i</sup>               | 87.17 (5)   | C3—C4—C5  | 120.02 (16) |
| O1W—Cd1—O1W <sup>i</sup>              | 180.00 (5)  | C3—C4—H4A | 120.0       |
| O3 <sup>i</sup> —Cd1—O2 <sup>i</sup>  | 81.00 (5)   | C5—C4—H4A | 120.0       |
| O3—Cd1—O2 <sup>i</sup>                | 99.00 (5)   | C6—C5—C4  | 118.38 (15) |
| O1W—Cd1—O2 <sup>i</sup>               | 91.77 (5)   | C6—C5—C7  | 118.06 (14) |

|                                       |              |               |              |
|---------------------------------------|--------------|---------------|--------------|
| O1W <sup>i</sup> —Cd1—O2 <sup>i</sup> | 88.23 (5)    | C4—C5—C7      | 123.54 (15)  |
| O3 <sup>i</sup> —Cd1—O2               | 99.00 (5)    | O1—C6—C5      | 120.21 (14)  |
| O3—Cd1—O2                             | 81.00 (5)    | O1—C6—C1      | 117.23 (15)  |
| O1W—Cd1—O2                            | 88.23 (5)    | C5—C6—C1      | 122.57 (15)  |
| O1W <sup>i</sup> —Cd1—O2              | 91.77 (5)    | C8—C7—C5      | 121.68 (15)  |
| O2 <sup>i</sup> —Cd1—O2               | 180.00 (3)   | C8—C7—H7A     | 119.2        |
| C9—O1—C6                              | 122.72 (13)  | C5—C7—H7A     | 119.2        |
| Cd1—O1W—H1WA                          | 100.6        | C7—C8—C9      | 119.32 (14)  |
| Cd1—O1W—H1WB                          | 100.5        | C7—C8—C10     | 118.64 (14)  |
| H1WA—O1W—H1WB                         | 130.4        | C9—C8—C10     | 122.04 (14)  |
| C9—O2—Cd1                             | 123.19 (11)  | O2—C9—O1      | 114.46 (14)  |
| C10—O3—Cd1                            | 132.88 (11)  | O2—C9—C8      | 127.61 (15)  |
| C6—C1—C2                              | 117.96 (16)  | O1—C9—C8      | 117.92 (14)  |
| C6—C1—H1A                             | 121.0        | O4—C10—O3     | 124.02 (15)  |
| C2—C1—H1A                             | 121.0        | O4—C10—C8     | 115.83 (14)  |
| C1—C2—C3                              | 120.77 (15)  | O3—C10—C8     | 120.11 (14)  |
| C1—C2—H2A                             | 119.6        |               |              |
| O3 <sup>i</sup> —Cd1—O2—C9            | 151.16 (12)  | C2—C1—C6—C5   | 0.0 (2)      |
| O3—Cd1—O2—C9                          | −28.84 (12)  | C6—C5—C7—C8   | −0.7 (2)     |
| O1W—Cd1—O2—C9                         | 64.31 (13)   | C4—C5—C7—C8   | −178.75 (15) |
| O1W <sup>i</sup> —Cd1—O2—C9           | −115.69 (13) | C5—C7—C8—C9   | −2.1 (2)     |
| O1W—Cd1—O3—C10                        | −87.46 (15)  | C5—C7—C8—C10  | 178.98 (14)  |
| O1W <sup>i</sup> —Cd1—O3—C10          | 92.54 (15)   | Cd1—O2—C9—O1  | −149.14 (10) |
| O2 <sup>i</sup> —Cd1—O3—C10           | −179.70 (15) | Cd1—O2—C9—C8  | 32.0 (2)     |
| O2—Cd1—O3—C10                         | 0.30 (15)    | C6—O1—C9—O2   | 179.45 (13)  |
| C6—C1—C2—C3                           | 0.0 (2)      | C6—O1—C9—C8   | −1.5 (2)     |
| C1—C2—C3—C4                           | −0.4 (3)     | C7—C8—C9—O2   | −177.95 (16) |
| C2—C3—C4—C5                           | 0.7 (2)      | C10—C8—C9—O2  | 0.9 (3)      |
| C3—C4—C5—C6                           | −0.7 (2)     | C7—C8—C9—O1   | 3.2 (2)      |
| C3—C4—C5—C7                           | 177.35 (15)  | C10—C8—C9—O1  | −177.92 (13) |
| C9—O1—C6—C5                           | −1.3 (2)     | Cd1—O3—C10—O4 | −156.56 (12) |
| C9—O1—C6—C1                           | 178.95 (14)  | Cd1—O3—C10—C8 | 25.5 (2)     |
| C4—C5—C6—O1                           | −179.43 (14) | C7—C8—C10—O4  | −31.7 (2)    |
| C7—C5—C6—O1                           | 2.4 (2)      | C9—C8—C10—O4  | 149.44 (15)  |
| C4—C5—C6—C1                           | 0.3 (2)      | C7—C8—C10—O3  | 146.42 (16)  |
| C7—C5—C6—C1                           | −177.83 (14) | C9—C8—C10—O3  | −32.5 (2)    |
| C2—C1—C6—O1                           | 179.78 (14)  |               |              |

Symmetry codes: (i)  $-x+2, -y+1, -z$ .

### Hydrogen-bond geometry (Å, °)

| $D-H\cdots A$                       | $D-H$ | $H\cdots A$ | $D\cdots A$ | $D-H\cdots A$ |
|-------------------------------------|-------|-------------|-------------|---------------|
| O1W—H1WA $\cdots$ O4 <sup>ii</sup>  | 0.85  | 1.90        | 2.6877 (18) | 153           |
| O1W—H1WB $\cdots$ O4 <sup>iii</sup> | 0.85  | 1.94        | 2.721 (2)   | 153           |

Symmetry codes: (ii)  $-x+1, -y+1, -z$ ; (iii)  $x+1, y+1, z$ .

Fig. 1




Fig. 2

